

KUMMER STRUCTURES

ADAM CHALCRAFT
MICHAEL FRYERS

Suppose we take an abelian group G and let $G/\pm 1$ be the quotient of G by the action of negation. What structure does $G/\pm 1$ inherit from the group structure of G ?

Let us write $\bar{x} \in G/\pm 1$ for the image of $x \in G$. Then given $\bar{x}, \bar{y} \in G/\pm 1$, we cannot define $\overline{x+y}$ uniquely, because $\overline{-x} = \bar{x}$ but $\overline{-x+y} \neq \overline{x+y}$ in general; but we can define the unordered pair $\{\overline{x+y}, \overline{x-y}\}$. We thus get a map κ from $(G/\pm 1)^{\{2\}}$, the set of unordered pairs of elements of $G/\pm 1$, to itself. We call the structure $(G/\pm 1, \kappa)$ the *Kummer* of G .

An example from geometry explains our use of the name *Kummer*: the quotient of an Abelian surface (i.e., a two-dimensional projective algebraic group) by ± 1 is called a *Kummer surface*.

In this paper we propose some axioms that hold for the structure $(G/\pm 1, \kappa)$, and show that every structure satisfying those axioms either is the Kummer of a unique group, or comes from one other construction, the quotient of a 2-torsion group by an involution. The proofs are constructive, showing how G can be reconstructed from $(G/\pm 1, \kappa)$.

1. AXIOMS

Definition. A *Kummer structure* is a set K with a map $\kappa: K^{\{2\}} \rightarrow K^{\{2\}}$ satisfying the following axioms: (we use the notation $a \ b \rightarrow c \ d$ to mean $\{a, b\} \xrightarrow{\kappa} \{c, d\}$)

- A1. There is an element $0 \in K$ such that for every $a \in K$ we have $a \ 0 \rightarrow a \ a$.
- A2. For every $a \in K$ there is an element $2a \in K$ such that $a \ a \rightarrow 0 \ 2a$.
- A3. Every $a, b, c \in K$ fit into a diagram like this:

$$\begin{array}{ccccc} & c & & c & c \\ a & b & \rightarrow & p_0 & p_1 \\ & \downarrow & & \downarrow & \downarrow \\ a & q_0 & \rightarrow & s_0 & s_1 \\ a & q_1 & \rightarrow & s_2 & s_3 \end{array}$$

(here the downward arrows mean $c \ b \rightarrow q_0 \ q_1$ etc.).

- A4. 2 is a homomorphism: that is, if $a \ b \rightarrow c \ d$ then $2a \ 2b \rightarrow 2c \ 2d$.

Remark. It is clear that the element 0 and the map 2 described in the axioms are uniquely determined by κ .

We will write $a \ b \rightarrow c \ \star$ to mean that $a \ b \rightarrow c \ d$ for some unspecified d , and use the obvious notation $4a := 2(2a)$.

2. EXAMPLES

Definition. Given an abelian group $(G, +)$, the *Kummer* of G is $(G/\pm 1, \kappa)$, where

$$G/\pm 1 := \{ \{x, -x\} \mid x \in G \},$$

and (writing $\overline{x}$ for $\{x, -x\}$) κ is defined by

$$\overline{x} \ \overline{y} \rightarrow \overline{x+y} \ \overline{x-y}.$$

Remark. It is easy to check that $G/\pm 1$ satisfies axioms A1–A4 with $0 = \overline{0}$ and $2\overline{x} = \overline{2x}$.

Example. Let K be the closed interval $[-1, 1]$, and let κ be defined by the rule

$$a \ b \rightarrow c \star \iff a^2 + b^2 + c^2 = 2abc + 1$$

(so $\kappa\{a, b\}$ is the set of solutions of this equation.) Then the axioms A1–A4 can be checked by hand. Alternatively, one may observe that this K is the Kummer of the group $\mathbb{R}/2\pi\mathbb{Z}$, where $\overline{\theta}$ is represented by $\cos \theta \in [-1, 1]$.

Example. (For algebraic geometers) If $G = E(k)$ is the set of points of an elliptic curve in Weierstraß form over an algebraically closed field k , then $K = G/\pm 1$ can be identified with $\mathbb{P}^1(k)$ by representing a point $\overline{(x, y)}$ by x . The map κ can be given by an algebraic formula like that in the previous example. If k is not algebraically closed, then K is the subset of $\mathbb{P}^1(k)$ consisting of x -coordinates of k -points of E .

Since the axioms were intended to capture the structure of Kummers of abelian groups, it would not be surprising if this were the only example of a Kummer structure. There is, however, another construction, which we will later show leads to Kummer structures not isomorphic to the Kummers of groups:

Definition. Let G be a 2-torsion abelian group, and $\iota: G \rightarrow G$ an involution (that is, an automorphism with $\iota^2 = 1$). The *twisted Kummer* of (G, ι) is $(G/\iota, \kappa)$, where

$$G/\iota := \{ \{x, \iota x\} \mid x \in G \},$$

and (writing $\overline{x}$ for $\{x, \iota x\}$) κ is defined by

$$\overline{x} \ \overline{y} \rightarrow \overline{x+y} \ \overline{x+\iota y}.$$

For convenience, we define a *twisted group* to be a 2-torsion abelian group together with an involution.

Remark. Again, it is easy to check the axioms, and that $0 = \overline{0}$ and $2\overline{x} = \overline{x + \iota x}$.

This construction still works in the special case where ι is the identity, but since, for a 2-torsion group, -1 is also the identity, we get nothing new in this case: $G/1 = G/\pm 1 = G$.

We shall show that every Kummer structure is either a Kummer or a twisted Kummer, and describe the few which have both constructions.

Example. We give one example of a structure which satisfies A1–A3 but not A4, to show that A4 is not redundant.

Let $Q_8 = \{1, -1, i, -i, j, -j, k, -k\}$ be the quaternion group of order 8. Quotienting Q_8 by the inverse operation gives a set of 5 elements, $K = \{\overline{1}, \overline{-1}, \overline{i}, \overline{j}, \overline{k}\}$. It is straightforward to check that we can define

$$\overline{x} \ \overline{y} \rightarrow \overline{xy} \ \overline{xy^{-1}}, \quad 0 = \overline{1}, \quad 2\overline{x} = \overline{x^2},$$

and the axioms A1–A3 then hold, but A4 fails: $\overline{i} \ \overline{j} \rightarrow \overline{k} \ \overline{k}$ should imply $\overline{-1} \ \overline{-1} \rightarrow \overline{-1} \ \overline{-1}$, but in fact $\overline{-1} \ \overline{-1} \rightarrow \overline{1} \ \overline{1}$.

(We can regard K as the ‘Kummer’ of Q_8 – but note that for most non-abelian groups we cannot even give a consistent definition for κ of this form: to be consistent we need $\kappa\{\overline{x}, \overline{y}\} = \kappa\{\overline{y}, \overline{x}\}$ to hold.)

3. FIRST PROPERTIES, AND THE 2-TORSION GROUP

In this section K shall be a Kummer structure. We will collect some useful consequences of the axioms, which lead us to a description of the 2-torsion elements of K – i.e., elements in the kernel of the map 2.

Lemma 1. *The following hold for any $a, b, c, d \in K$.*

- L1. $20 = 0$.
- L2. $a b \rightarrow c \star \Leftrightarrow b c \rightarrow a \star$.
- L3. $a b \rightarrow 0 \star \Leftrightarrow a = b$.
- L4. $a b \rightarrow c d \Rightarrow c d \rightarrow 2a \ 2b$.
- L5. Let $a b \rightarrow c d$. Then $c = d \Leftrightarrow (2a = 0 \text{ or } 2b = 0)$.
- L6. If $2a = 2b = 0$ then $a b \rightarrow c c$ for some c with $2c = 0$.
- L7. Given the notation of A3, there are also r_0, r_1 such that

$$\begin{array}{c} c \\ a \\ \downarrow \\ b \ r_0 \rightarrow s_0 \ s_3 \\ b \ r_1 \rightarrow s_2 \ s_1. \end{array}$$

Proof. L1. A1 $\Rightarrow 0 \ 0 \rightarrow 0 \ 0$; but A2 $\Rightarrow 0 \ 0 \rightarrow 0 \ 20$; so $20 = 0$.

L2. Given $a b \rightarrow c \star$, $b b \rightarrow 0 \ 2b$ and $a \ 0 \rightarrow a \ a$ are axioms, so we can fill in this case of A3:

$$\begin{array}{ccc} b & & b \ b \\ a \ b & \rightarrow & c \ \star \\ \downarrow & & \downarrow \downarrow \\ a \ 0 & \rightarrow & a \ a \\ a \ 2b & \rightarrow & \star \ \star, \end{array}$$

and read off $b c \rightarrow a \star$. The other implication follows by symmetry.

L3. Put $c = 0$ in L2: by A1, $b \ 0 \rightarrow a \star \Leftrightarrow a = b$.

L4. We first prove $c d \rightarrow 2a \star$: we have (using A3 and L2)

$$\begin{array}{ccc} c & & c \ c \\ a \ b & \rightarrow & c \ d \\ \downarrow & & \downarrow \downarrow \\ a \ a & \rightarrow & p \ q \\ a \ e & \rightarrow & r \ s, \end{array}$$

and A2 implies $\{p, q\} = \{0, 2a\}$. If $q = 2a$ then we read off $c d \rightarrow 2a \star$. If $p = 2a$ then we read off $c c \rightarrow 2a \star$, but also $c d \rightarrow 0 \star$, so that (by L3) $c = d$, so again $c d \rightarrow 2a \star$.

Symmetry gives also $c d \rightarrow 2b \star$, so we're done unless $2a = 2b$.

By A4, $2a \ 2b \rightarrow 2c \ 2d$, and if $2a = 2b$ then A2 implies that one of $2c$ and $2d$ is 0. The two cases are the same, so take $2c = 0$. Now we can complete the diagram above: $p = r = 0$ by A2; $e = a$ by L3; $q = s = 2a = 2b$ by A2; and we're done.

L5. L4 says $a b \rightarrow c d \Rightarrow c d \rightarrow 2a \ 2b$; so, by L3, $c = d \Leftrightarrow (2a = 0 \text{ or } 2b = 0)$.

L6. L5 $\Rightarrow a b \rightarrow c c$ for some c ; by A4, $2a \ 2b \rightarrow 2c \ 2c$; but $\{2a, 2b\} = \{0, 0\}$ and $0 \ 0 \rightarrow 0 \ 0$, so $2c = 0$.

L7. Three applications of A3 give

$$\begin{array}{ccccc}
 & c & & c & c \\
 a & b & \rightarrow & p_0 & p_1 \\
 & \downarrow & & \downarrow & \downarrow \\
 a & q_0 & \rightarrow & s_0 & s_1 \\
 a & q_1 & \rightarrow & s_2 & s_3,
 \end{array}
 \quad
 \begin{array}{ccccc}
 & c & & c & c \\
 b & a & \rightarrow & p_0 & p_1 \\
 & \downarrow & & \downarrow & \downarrow \\
 b & r_0 & \rightarrow & t_0 & t_1 \\
 b & r_1 & \rightarrow & t_2 & t_3,
 \end{array}
 \quad
 \begin{array}{ccccc}
 & b & & b & b \\
 a & c & \rightarrow & r_0 & r_1 \\
 & \downarrow & & \downarrow & \downarrow \\
 a & q_0 & \rightarrow & u_0 & u_1 \\
 a & q_1 & \rightarrow & u_2 & u_3.
 \end{array}$$

Switching r_0 and r_1 if necessary, we may assume $t_0 = s_0$ and $t_2 = s_2$. If $t_1 = s_3$ and $t_3 = s_1$, we have L7, so assume $t_1 = s_1 \neq t_3 = s_3$. If $s_0 = s_2$, we can switch r_0 and r_1 to give L7, so assume also $s_0 \neq s_2$. We will now derive a contradiction.

From the third diagram, $u_1 \in \{s_0, s_1\} \cap \{t_2, t_3\}$, so either $u_1 = s_0 = s_3$ or $u_1 = s_1 = s_2$. Switching p_0 and p_1 if necessary, we may assume $u_1 = s_1 = s_2$, and then $s_i = t_i = u_i$ for each i .

Now (by L4)

$$\begin{aligned}
 a \quad q_0 &\rightarrow s_0 \quad s_1 \rightarrow 2a \quad 2q_0, \\
 b \quad r_0 &\rightarrow s_0 \quad s_1 \rightarrow 2b \quad 2r_0, \\
 c \quad p_0 &\rightarrow s_0 \quad s_1 \rightarrow 2c \quad 2p_0,
 \end{aligned}$$

so two out of $2a$, $2b$, and $2c$ are equal; without loss of generality $2a = 2b$. But

$$a \quad b \rightarrow q_0 \quad q_1 \rightarrow 2a \quad 2b,$$

so by L5, $2q_0 = 0$ or $2q_1 = 0$. By L5 again, if $2q_0 = 0$ then $s_0 = s_1$, but we assumed $s_0 \neq s_2 = s_1$; if $2q_1 = 0$ then $s_2 = s_3$, but we assumed $s_2 = s_1 \neq s_3$. This is our contradiction. $\square$

Definition. The 2-torsion of K is the set $K[2] := \{a \in K \mid 2a = 0\}$. By L6, if $a, b \in K[2]$ then $a \quad b \rightarrow c \quad c$ for some $c \in K[2]$; we define $a + b = c$ in this case.

Lemma 2. This construction makes $K[2]$ into a 2-torsion abelian group. For an abelian group G we have $(G/\pm 1)[2] \cong G[2]$, the 2-torsion in G . For a twisted group (G, ι) , we have $(G/\iota)[2] \cong G[1 + \iota] := \ker(1 + \iota) \subseteq G$.

Proof. Clearly $+$ is commutative. For any $a, b, c \in K[2]$, A3 gives us a diagram

$$\begin{array}{ccccc}
 & c & & c & c \\
 a & b & \rightarrow & (a+b) & (a+b) \\
 & \downarrow & & \downarrow & \downarrow \\
 a & (b+c) & \rightarrow & d & d \\
 a & (b+c) & \rightarrow & d & d,
 \end{array}$$

so $a + (b + c) = d = (a + b) + c$; that is, $+$ is associative. For any $a \in K[2]$ we have $a \quad a \rightarrow 0 \quad 0$ by A2, so $a + a = 0$. Finally A1 gives us $a + 0 = a$. So $K[2]$ is a 2-torsion abelian group.

For G an abelian group and $x, y \in G[2]$ we have $\overline{x \quad y} \rightarrow \overline{x+y} \quad \overline{x+y}$, so $\overline{x+y} = \overline{x+y}$; that is, $(x \mapsto \overline{x})$ is a group homomorphism $G[2] \rightarrow (G/\pm 1)[2]$. It is clearly bijective, and so an isomorphism.

Exactly the same argument applies to (G, ι) a twisted group and $x, y \in G[1 + \iota]$. $\square$

Remark. In particular, this shows that 2-torsion Kummer structures are essentially the same as 2-torsion abelian groups (or as twisted groups with $\iota = 1$).

4. STRINGS

Since the previous section completely analyses 2-torsion Kummer structures, from now on we assume that we have some non-2-torsion element: in this section K shall be a Kummer structure and $g \in K \setminus K[2]$.

Definition. Define $K_g \subseteq K^{\mathbb{Z}}$ to be the set of sequences $\alpha = (\alpha_n)_{n \in \mathbb{Z}}$ such that $\alpha_n g \rightarrow \alpha_{n-1} \alpha_{n+1}$ for all n . We call elements of K_g *strings*.

Lemma 3. *For any $n \in \mathbb{Z}$, any $\alpha_n, \alpha_{n+1} \in K$ such that $\alpha_n g \rightarrow \alpha_{n+1} \star$ can be extended to a unique $\alpha \in K_g$.*

Proof. L2 says that the condition $\alpha_n g \rightarrow \alpha_{n+1} \star$ is symmetric in α_n and α_{n+1} .

Given such α_n and α_{n+1} , there is a unique α_{n-1} such that $\alpha_n g \rightarrow \alpha_{n+1} \alpha_{n-1}$ and a unique α_{n+2} such that $\alpha_{n+1} g \rightarrow \alpha_n \alpha_{n+2}$.

By induction on n in both directions, this construction determines α_m for all $m \in \mathbb{Z}$. $\square$

Lemma 4. *If $\alpha \in K_g$ then $\alpha_n 2g \rightarrow \alpha_{n-2} \alpha_{n+2}$ for any n .*

Proof. By A3, A2, and the definition of strings,

$$\begin{array}{ccccc} & g & & g & g \\ \alpha_n & g & \rightarrow & \alpha_{n-1} & \alpha_{n+1} \\ & \downarrow & & \downarrow & \downarrow \\ \alpha_n & 0 & \rightarrow & \alpha_n & \alpha_n \\ \alpha_n & 2g & \rightarrow & \alpha_{n-2} & \alpha_{n+2}. \end{array}$$

$\square$

Definition. Define $\circ$ to be the unique element of K_g such that $\circ_0 = 0$. (Its existence and uniqueness are guaranteed by Lemma 3: since $0 g \rightarrow g g$, the only possibility for $\circ_1$ is g .)

If $\alpha \in K_g$, define $\rho(\alpha)_n := \alpha_{-n}$; clearly also $\rho(\alpha) \in K_g$.

If $\alpha, \beta, \gamma, \delta \in K_g$ and for all $n, m \in \mathbb{Z}$,

$$\alpha_n \beta_m \rightarrow \gamma_{n+m} \delta_{n-m},$$

then we write $\gamma = \alpha + \beta$ and $\delta = \alpha - \beta$.

If $\alpha, \beta, \tilde{\gamma}, \tilde{\delta} \in K_g$ and for all $n, m \in \mathbb{Z}$,

$$\begin{array}{ll} \alpha_n \beta_m \rightarrow \tilde{\gamma}_{n+m} \tilde{\delta}_{n-m} & \text{if } n \equiv m \pmod{2}, \\ \alpha_n \beta_m \rightarrow \tilde{\gamma}_{n-m} \tilde{\delta}_{n+m} & \text{if } n \not\equiv m \pmod{2}, \end{array}$$

then we write $\gamma = \alpha \oplus \beta$ and $\delta = \alpha \ominus \beta$.

Remark. In Lemma 12 of section 5 we shall justify the functional notation by showing that these definitions make $+$, $-$, $\oplus$, and $\ominus$ into partial functions $K_g \times K_g \rightarrow K_g$. (i.e., for any α, β there is at most one γ such that $\gamma = \alpha + \beta$, etc..)

For now, note that $\delta = \alpha - \beta \Leftrightarrow \delta = \alpha + \rho(\beta)$, and $\tilde{\delta} = \alpha \ominus \beta \Leftrightarrow \tilde{\delta} = \alpha \oplus \rho(\beta)$.

Theorem 5. *Assume Lemma 12, that $+$ and $\oplus$ are partial functions. Let K be a Kummer structure and $g \in K \setminus K[2]$.*

$K \cong G/\pm 1$ for an abelian group G if and only if taking $+$ as addition, $\circ$ as zero, and ρ as negation makes K_g into a group isomorphic to G .

$K \cong G/\iota$ for a twisted group (G, ι) if and only if taking $\oplus$ as addition, $\circ$ as zero, and ρ as the involution makes K_g into a twisted group isomorphic to (G, ι) .

Proof. If $(+, \circ, \rho)$ make K_g into an abelian group, the map $K_g \rightarrow K$ taking $\alpha \mapsto \alpha_0$ clearly factors through a map $K_g/\pm 1 = K_g/\rho \rightarrow K$; Lemma 3 implies that this map is a bijection $K_g/\pm 1 \leftrightarrow K$, and it is then straightforward to check that it is an isomorphism of Kummer structures.

On the other hand, if $K = G/\pm 1$, we have $g = \bar{z} = \{z, -z\}$ for some $z \in G \setminus G[2]$. Define a map $\phi: G \rightarrow K_g$ by $\phi(x)_n = \overline{x + nz}$, which clearly makes $\phi(x)$ a $\bar{z}$ -string. Now ϕ is surjective, since for any $\alpha \in K_g$ with $\alpha_0 = \bar{x}$, either $\alpha_1 = \overline{x + z}$, in which case (by Lemma 3) $\alpha = \phi(x)$, or $\alpha_1 = \overline{x - z}$, in which case $\alpha = \phi(-x)$. And ϕ is injective, for if $\phi(y) = \phi(x)$ then $\bar{y} = \bar{x}$ and $\overline{y + z} = \overline{x + z}$, so $y = \pm x$ and $y + z = \pm(x + z)$; but these imply either $y = x$ or $2z = 0$, and the latter we know is false.

It is straightforward to check from the definitions above that $\phi(x + y) = \phi(x) + \phi(y)$ and $\phi(x - y) = \phi(x) - \phi(y)$, so that $+$ is an associative total function, and that $\phi(-x) = \rho(\phi(x))$ and $\phi(0) = 0_{K_g}$, so that K_g is an abelian group as claimed and $\phi: G \cong K_g$.

The proof of the twisted group version of this result is similar, and we omit it. $\square$

Remark. The ideas of this section may also be used to define a natural action of the multiplicative monoid of $\mathbb{N}$ on K : take $0a = 0$, $1a = a$, and the rule

$$na \quad a \rightarrow (n-1)a \quad (n+1)a \quad (n > 0).$$

Clearly $2a$ so defined is the same as the $2a$ we have been using, since A2 is a special case of the rule above. We do not make use of this construction, so we leave the interested reader to check that this is well-defined, and that this definition makes each $n \in \mathbb{N}$ into a homomorphism in the sense of axiom A4.

5. COLOURING DIAMOND GRIDS

In this section, K shall be a Kummer structure, $g \in K \setminus K[2]$, and α, β shall be two elements of K_g . We wish to explore the question of existence and uniqueness of $\alpha + \beta$, $\alpha - \beta$, $\alpha \oplus \beta$, and $\alpha \ominus \beta$.

Definition. Let D be the graph whose nodes are pairs $(n, m) \in \mathbb{Z}^2$, with (n, m) adjacent to (n', m') if and only if $|n - n'| = |m - m'| = 1$. The edges of this graph lie in two directions: a *rising* edge connects (n, m) and $(n + 1, m + 1)$; a *falling* edge connects (n, m) and $(n + 1, m - 1)$. For each $p \in \{0, 1\}$ let D_p be the component of D whose nodes are (n, m) with $n + m \equiv p \pmod{2}$.

If $\alpha_n \beta_m \rightarrow a \ b$, then call a, b the *values* at the node (n, m) . When we draw parts of D or D_p , we'll simply label each node with its unordered pair of values.

Lemma 6. (The diamond rule) *Every diamond of nodes of D_p looks like*

$$\begin{array}{ccc} & c \ b & \\ & \swarrow \quad \searrow & \\ a \ b & & c \ d, \\ & \nwarrow \quad \nearrow & \\ & a \ d & \end{array}$$

for some $a, b, c, d \in K$ (not necessarily distinct).

Proof. In other words, for any $n, m \in \mathbb{Z}$ (with $n + m \not\equiv p \pmod{2}$), there are $a, b, c, d \in K$ such that

$$\begin{array}{ccc} & \alpha_n \beta_{m+1} \rightarrow c \ b & \\ \alpha_{n-1} \beta_m \rightarrow a \ b & & \alpha_{n+1} \beta_m \rightarrow c \ d. \\ & \alpha_n \beta_{m-1} \rightarrow a \ d & \end{array}$$

But this is simply a case of A3:

$$\begin{array}{ccccc}
 & \alpha_n & & \alpha_n & \alpha_n \\
 \beta_m & g & \rightarrow & \beta_{m-1} & \beta_{m+1} \\
 & \downarrow & & \downarrow & \downarrow \\
 \beta_m & \alpha_{n-1} & \rightarrow & a & b \\
 \beta_m & \alpha_{n+1} & \rightarrow & d & c .
 \end{array}$$

□

Lemma 7. *The node of D_{1-p} in the middle of the diamond of Lemma 6 has values $u v$, where $u g \rightarrow a c$ and $v g \rightarrow b d$.*

Proof. L7, applied to the instance of A3 in the proof of Lemma 6, states that $\alpha_n \beta_n \rightarrow u v$ such that $u g \rightarrow a c$ and $v g \rightarrow b d$. □

Lemma 8. (The linear rule) *For any straight-line path in D , all the nodes in it have a value in common.*

Proof. From the diamond rule we see that any two adjacent nodes have a value in common, so in any exception to the lemma the nodes must contain a subsequence of the form $a c - b c - \dots - b c - b d$, where $a \neq b \neq c \neq d$, and $b c$ is repeated $n \geq 1$ times. Let us suppose we have such a path in D , with n minimal.

Pick one of the two adjacent parallel paths. Applying the diamond rule to the diamonds between the two paths, in turn starting from the left, the values on the neighbouring path are $a \star - b \star - \dots - b \star - b \star$. Now, starting from the right and applying the diamond rule again, we can fill in the remaining values: the neighbouring path also has values $a c - b c - \dots - b c - b d$. So, applying Lemma 7, we have a part of D that looks (up to a possible 90° rotation) like

$$\begin{array}{ccc}
 & a c & \\
 a c & \swarrow \quad \searrow & \\
 u_0 v_0 & & b c \\
 & \swarrow \quad \searrow & \\
 b c & u_1 v_1 & \dots \\
 & \swarrow \quad \searrow & \\
 & \dots & b c \\
 & \swarrow \quad \searrow & \\
 b c & u_n v_n & b d, \\
 & \swarrow \quad \searrow & \\
 & b d &
 \end{array}
 \quad
 \begin{array}{l}
 u_0 g \rightarrow c c, \\
 v_0 g \rightarrow a b, \\
 \\
 u_i g \rightarrow c c, \\
 v_i g \rightarrow b b, \\
 (0 < i < n) \\
 \\
 u_n g \rightarrow c d, \\
 v_n g \rightarrow b b.
 \end{array}$$

Since the path from $u_0 v_0$ to $u_n v_n$ is shorter than our minimal bad path, there must be a value in common between $\{u_0, v_0\}, \dots, \{u_n, v_n\}$. Since $a \neq b \neq c \neq d$, the conditions on the right above make this impossible unless $n = 1$, and then $v_0 \neq v_1 \neq u_0 \neq u_1$, so the common value must be $v_0 = u_1$, which implies $c = a$ and $d = b$. Renaming $u := u_0$, $v := v_0 = u_1$, and $w := v_1$, with one more application of the diamond rule we have

$$\begin{array}{ccc}
 & a a & u \star \\
 a a & \swarrow \quad \searrow & \\
 u v & & b a \\
 & \swarrow \quad \searrow & \\
 b a & w v & b b, \\
 & \swarrow \quad \searrow & \\
 & b b &
 \end{array}
 \quad
 \begin{array}{l}
 u g \rightarrow a a, \\
 v g \rightarrow a b, \\
 w g \rightarrow b b.
 \end{array}$$

By Lemma 7, either $bg \rightarrow uw$ or $ag \rightarrow uw$. But by L2, since $vg \rightarrow ab$, both $ag \rightarrow v\star$ and $bg \rightarrow v\star$. This is impossible, since we know $u \neq v \neq w$. $\square$

Definition. To *colour* a node N shall mean to assign values Γ_N and $\Delta_N \in K$ such that $\{\Gamma_N, \Delta_N\}$ is the value set at N . A colouring of all the nodes of D_p shall be said to be *consistent* if whenever AB is a falling edge, $\Gamma_A = \Gamma_B$, and whenever AB is a rising edge, $\Delta_A = \Delta_B$.

Lemma 9. D_p has a consistent colouring, unique unless every node of D_p has the same value set and that common value set is doubleton, in which case D_p has two consistent colourings.

Proof. Say that an edge of D_p is *even* if the value sets of the two nodes it joins are equal. Looking at the diamond of Lemma 6, the upper left edge is even if and only if $a = c$, if and only if the lower right edge is even. Similarly the upper right and lower left edges are each even if and only if $b = d$. Thus even edges occur in infinite *ladders* of parallel edges.

Let D'_p be the graph obtained by contracting all even edges of D_p . Because only even edges have been contracted, the nodes of D'_p inherit well-defined value sets from the nodes of D_p . Because even edges form ladders, the underlying graph of D'_p is isomorphic to a subgraph of D_p consisting of the nodes $\{(n, m) \in D_p \mid \lambda_+ \leq n + m \leq \mu_+ \text{ and } \lambda_- \leq n - m \leq \mu_-\}$, for some $-\infty \leq \lambda_{\pm} \leq \mu_{\pm} \leq \infty$. Clearly the notion of rising and falling edges can be applied naturally to D'_p , and the diamond rule and the linear rule for D'_p follow from the same for D_p . For an even edge AB , whether rising or falling, the condition that a colouring be consistent is equivalent to $\Gamma_A = \Gamma_B$ and $\Delta_A = \Delta_B$, so consistent colourings of D_p and of D'_p are in one-to-one correspondence.

D'_p has no even edges, so given any edge AB of D'_p , there is only one value in common between A and B , so there is only one possible consistent colouring of just A and B . (If the edge is falling, the common value must be $\Gamma_A = \Gamma_B$, if rising, $\Delta_A = \Delta_B$.) Say that the edge AB *forces* those particular colourings of A and B .

If there is at least one non-even edge in D_p , then every vertex A of D'_p meets an edge AB , which forces a particular colouring of A . So there is at most one consistent colouring of D'_p , and so of D_p .

In this case, to prove the existence of a consistent colouring of D'_p , we must show that any two coincident edges AB, BC force the same colouring on their common end B .


But if AB and BC are not parallel, then they are part of a diamond, and the diamond rule states that any diamond can be consistently coloured. If AB and BC are parallel, they are a straight path, and the linear rule states that such a path can be consistently coloured.

So AB and BC must force the same colouring on B .

On the other hand, if every edge of D_p is even, which is to say the value sets at all nodes of D_p are equal, D'_p is a single node, and any colouring of a single node is consistent; so there are two consistent colourings if the value set is doubleton and just one if the value set is singleton. $\square$

Lemma 10. At most one of the components D_0 and D_1 has two distinct consistent colourings.

Proof. Suppose we have a counterexample. Then by Lemma 9 all value sets at nodes of D_p are equal, for each p . So each diamond of D_0 looks like



Since $2g \neq 0$, by L5 we must have $2c = 2d = 0$. Similarly $2a = 2b = 0$. But by A4, if $c \rightarrow a$ then $2c \rightarrow 2a$ and $2a = 0$. So, by L3, we have $2g = 2c = 0$, a contradiction. $\square$

Now we can rephrase the definitions of $+$, $-$, $\oplus$, and $\ominus$ from Section 4 in terms of colourings of D :

Lemma 11. For $\gamma, \delta, \tilde{\gamma}, \tilde{\delta} \in K_g$, respectively

$$\begin{aligned}\gamma &= \alpha + \beta; \\ \delta &= \alpha - \beta; \\ \tilde{\gamma} &= \alpha \oplus \beta; \\ \tilde{\delta} &= \alpha \ominus \beta\end{aligned}$$

if and only if there is a consistent colouring $(\Gamma_{\cdot, \cdot}, \Delta_{\cdot, \cdot})$ of D such that, for all $n, m \in \mathbb{Z}$, respectively

$$\begin{aligned}\gamma_{n+m} &= \Gamma_{n,m}; \\ \delta_{n-m} &= \Delta_{n,m}; \\ \tilde{\gamma}_{n+m} &= \begin{cases} \Gamma_{n,m} & \text{if } n \equiv m \pmod{2}, \\ \Delta_{n,-m} & \text{if } n \not\equiv m \pmod{2}; \end{cases} \\ \tilde{\delta}_{n-m} &= \begin{cases} \Delta_{n,m} & \text{if } n \equiv m \pmod{2}, \\ \Gamma_{n,-m} & \text{if } n \not\equiv m \pmod{2}.\end{cases}\end{aligned}$$

$\square$

Lemma 12. There can be at most one string in K_g satisfying each of these conditions.

Proof. This is clear from Lemma 11 in the case where D has only one consistent colouring. In the remaining case, by Lemma 9 and Lemma 10, there are two consistent colourings $(\Gamma_{\cdot, \cdot}, \Delta_{\cdot, \cdot})$, $(\Gamma'_{\cdot, \cdot}, \Delta'_{\cdot, \cdot})$ of D : on one component of D , say D_p , the two are identical, while on the other, D_{1-p} , they satisfy $\Gamma' = \Delta$ and $\Delta' = \Gamma$. Lemma 9 further says that Γ and Γ' must each be constant on D_{1-p} , and not equal to each other.

From $(\Gamma_{\cdot, \cdot}, \Delta_{\cdot, \cdot})$ we can derive sequences $\gamma, \delta, \tilde{\gamma}, \tilde{\delta} \in K^{\mathbb{Z}}$ by the equations given in Lemma 11, and in the same way from $(\Gamma'_{\cdot, \cdot}, \Delta'_{\cdot, \cdot})$ we can derive $\gamma', \delta', \tilde{\gamma}', \tilde{\delta}' \in K^{\mathbb{Z}}$.

Suppose both γ and γ' are in K_g . Then $\gamma_p g \rightarrow \gamma_{p-1} \gamma_{p+1}$ and $\gamma'_p g \rightarrow \gamma'_{p-1} \gamma'_{p+1}$; in other words both $\Gamma_{p,0} g \rightarrow \Gamma_{p-1,0} \Gamma_{p+1,0}$ and $\Gamma'_{p,0} g \rightarrow \Gamma'_{p-1,0} \Gamma'_{p+1,0}$. But $\Gamma_{p,0} = \Gamma'_{p,0}$ whereas $\Gamma_{p-1,0} = \Gamma_{p+1,0} \neq \Gamma'_{p-1,0} = \Gamma'_{p+1,0}$, so this is impossible.

Similarly, at most one of each of the pairs $\{\delta, \delta'\}$, $\{\tilde{\gamma}, \tilde{\gamma}'\}$, and $\{\tilde{\delta}, \tilde{\delta}'\}$ can be in K_g . $\square$

Remark. This fulfils the promise made in Section 4, to prove that $+$, $-$, $\oplus$, and $\ominus$ are partial functions.

6. NON-4-TORSION KUMMER STRUCTURES

Recall that we defined $4g = 2(2g)$. In this section we consider the constructions of the previous two sections in the case $4g \neq 0$, and show that in this case K is the Kummer of a group. Throughout this section $r \equiv s$ always means modulo 2. First we need a quick lemma:

Lemma 13. If $\alpha \in K_g$ with $4g \neq 0$ then it cannot be that $\alpha_{n-2} = \alpha_n = \alpha_{n+2}$ for any n .

Proof. By Lemma 4, we would have $\alpha_n 2g \rightarrow \alpha_n \alpha_n$, so by L5, since $4g \neq 0$ we must have $2\alpha_n = 0$. But now applying A4 we have $2\alpha_{n+1} 2g \rightarrow 2\alpha_n 2\alpha_{n+2} = 0$, so by L3 we have $2\alpha_{n+1} = 2g$ and by L5, $4g = 0$ after all. $\square$

Theorem 14. *If K is a Kummer structure and there is any $g \in K$ with $4g \neq 0$ then $K \cong G/\pm 1$ for an abelian group G .*

Proof. By Theorem 5, the theorem will follow if we can show that $+$ and $-$ are total functions $K_g \times K_g \rightarrow K_g$, and that K_g is made into an abelian group by taking $+$ as addition, o as zero, and ρ as negation: then $G \cong K_g$.

First take $\alpha, \beta \in K_g$, construct the graph D as in Section 5 and choose a consistent colouring $(\Gamma_{\cdot, \cdot}, \Delta_{\cdot, \cdot})$ of it. Since the colouring is consistent, we can define $\gamma, \delta \in K^{\mathbb{Z}}$ by (for all $n, m \in \mathbb{Z}$)

$$\begin{aligned}\gamma_{n+m} &= \Gamma_{n,m}; \\ \delta_{n-m} &= \Delta_{n,m}.\end{aligned}$$

We need first to prove that $\gamma, \delta \in K_g$, so that we have $\gamma = \alpha + \beta$ and $\delta = \alpha - \beta$.

For any $r \equiv s$, we can write $r = n + m$, $s = n - m$ and apply Lemma 7 to the diamond centred at (n, m) to give

$$\begin{aligned}(1) \quad & \text{either } \begin{cases} \gamma_r \ g \rightarrow \gamma_{r-1} \ \gamma_{r+1}, \\ \delta_s \ g \rightarrow \delta_{s-1} \ \delta_{s+1}; \end{cases} \\ (2) \quad & \text{or } \begin{cases} \gamma_r \ g \rightarrow \delta_{s-1} \ \delta_{s+1}, \\ \delta_s \ g \rightarrow \gamma_{r-1} \ \gamma_{r+1}. \end{cases}\end{aligned}$$

Now $\gamma, \delta \in K_g \Leftrightarrow (1)$ holds for all r, s . So suppose there is some $r \equiv s \equiv p$ for which one of the equations of (1) fails. Then both equations of (2) hold, and so both equations of (1) fail. But one of these equations depends only on r and the other only on s , so (1) must fail and (2) hold for all $r \equiv s \equiv p$.

Suppose (1) fails for all $r, s \equiv p$ but holds for all $r, s \not\equiv p$. Then for any $r \equiv p$, we have (by L2)

$$\begin{aligned}\gamma_{r-1} \ g \rightarrow \gamma_r \ \star &\Leftrightarrow \gamma_r \ g \rightarrow \gamma_{r-1} \ \star, \\ \gamma_{r+1} \ g \rightarrow \gamma_r \ \star &\Leftrightarrow \gamma_r \ g \rightarrow \gamma_{r+1} \ \star,\end{aligned}$$

but $\gamma_r \ g \not\rightarrow \gamma_{r-1} \ \gamma_{r+1}$, so $\gamma_{r-1} = \gamma_{r+1}$ and (by L5) $2g = 0$ or $2\delta_r = 0$. The former is a contradiction, but given the latter for all $r \equiv p$, we can apply A4 to $\delta_{r+1} \ g \rightarrow \delta_r \ \delta_{r+2}$ to get $2\delta_{r+1} \ 2g \rightarrow 0 \ 0$, whence $4g = 0$, also a contradiction.

Therefore if (1) fails at all, it fails and (2) holds for all $r \equiv s$. An application of A3,

$$\begin{array}{ccc} g & g & g \\ \gamma_0 \ g \rightarrow \delta_{-1} \ \delta_1 & & \\ \downarrow & \downarrow & \downarrow \\ \gamma_0 \ 0 \rightarrow \gamma_0 \ \gamma_0 & & \\ \gamma_0 \ 2g \rightarrow \gamma_2 \ \gamma_2, & & \end{array}$$

gives $\gamma_0 \ 2g \rightarrow \gamma_2 \ \gamma_2$, so by L5, either $4g = 0$ (contradiction) or $2\gamma_0 = 0$. Similarly $2\gamma_2 = 0$. But then A4 applied to $\delta_1 \ g \rightarrow \gamma_0 \ \gamma_2$ gives $2\delta_1 \ 2g \rightarrow 0 \ 0$, so $4g = 0$, contradiction again.

So we always have $\gamma, \delta \in K_g$ and $+$ and $-$ are total functions. We must show that $(+, o, \rho)$ make K_g into an abelian group; that is, that $+$ is associative and commutative, that $\alpha + o = \alpha$ and that $\alpha + \rho(\alpha) = o$.

$+$ is obviously commutative.

By definition $\alpha_n \ o_0 \rightarrow (\alpha + o)_n \ (\alpha - o)_n$. But $o_0 = 0$, so $\alpha_n \ o_0 \rightarrow \alpha_n \ \alpha_n$. So $\alpha + o = \alpha$ ($= \alpha - o$).

By definition $\alpha_n \ \rho(\alpha)_{-n} \rightarrow (\alpha + \rho(\alpha))_0 \ (\alpha - \rho(\alpha))_{2n}$. But $\rho(\alpha)_{-n} = \alpha_n$, so $\alpha_n \ \rho(\alpha)_{-n} \rightarrow 0 \ \star$. So either $(\alpha + \rho(\alpha))_0 = 0$ or $(\alpha - \rho(\alpha))_{2n} = 0$. But the latter (for all n) contradicts Lemma 13, so $(\alpha + \rho(\alpha))_0 = 0$, which by definition of o means $\alpha + \rho(\alpha) = o$.

All that's left is to show $+$ to be associative. Fix $\alpha, \beta, \gamma \in K_g$, and set $\delta = \alpha + (\beta + \gamma)$ and $\delta' = (\alpha + \beta) + \gamma$.

Claim. For any n , if $\delta_n \neq \delta'_n$ then either $\delta_{n+2} = \delta_n$ or $\delta_{n+4} = \delta_n$, and either $\delta_{n-2} = \delta_n$ or $\delta_{n-4} = \delta_n$.

Proof. For any $p + q + r = n$, consider this instance of A3:

$$\begin{array}{ccc} \alpha_p & \begin{array}{c} \gamma_r \\ \beta_q \end{array} & \rightarrow (\alpha + \beta)_{p+q} \quad (\alpha - \beta)_{p-q} \\ & \downarrow & \downarrow \quad \downarrow \\ \alpha_p & (\beta + \gamma)_{q+r} & \rightarrow x \quad y \\ \alpha_p & (\beta - \gamma)_{q-r} & \rightarrow \star \quad \star \end{array}.$$

One of x and y is $(\alpha + (\beta + \gamma))_{p+q+r} = \delta_n$. Since $\delta_n \neq \delta'_n = ((\alpha + \beta) + \gamma)_{p+q+r}$, we must have

$$(3) \quad \delta_n = \text{one of } \begin{cases} ((\alpha + \beta) - \gamma)_{p+q-r} =: (\varepsilon^0)_{m_0}, \\ ((\alpha - \beta) + \gamma)_{p-q+r} =: (\varepsilon^1)_{m_1}, \\ ((\alpha - \beta) - \gamma)_{p-q-r} =: (\varepsilon^2)_{m_2}. \end{cases}$$

The constraint $p + q + r = n$ is equivalent to $m_0 \equiv m_1 \equiv m_2 \equiv n$ and $m_0 + m_1 - m_2 = n$.

For each i , we can't have $\delta_n = \varepsilon_{m_i}^i$ for every $m_i \equiv n$, by Lemma 13. So for any permutation $\{i, j, k\} = \{0, 1, 2\}$, we can find m_j and m_k such that $\varepsilon_{m_j}^j, \varepsilon_{m_k}^k \neq \delta_n$. Choosing m_i such that $m_0 + m_1 - m_2 = n$, by (3) we must have $\varepsilon_{m_i}^i = \delta_n$.

So each of the strings ε^i contains δ_n , so (by Lemma 3) each of these strings is obtained from δ by some translation and perhaps reversal. So if $\delta_{n+2}, \delta_{n+4} \neq \delta_n$, or $\delta_{n-4}, \delta_{n-2} \neq \delta_n$, we can find m_0 such that $\varepsilon_{m_0}^0, \varepsilon_{m_0+2}^0 \neq \delta_n$, and similarly m_1 such that $\varepsilon_{m_1}^1, \varepsilon_{m_1+2}^1 \neq \delta_n$. But then setting $m_2 = m_0 + m_1 - n$, by (3) we must have $\varepsilon_{m_2}^2 = \varepsilon_{m_2+2}^2 = \varepsilon_{m_2+4}^2 = \delta_n$, contradicting Lemma 13. This proves the Claim. $\square$

We can now complete the proof of Theorem 14. Suppose there is an n such that $\delta_n \neq \delta'_n$. Then we can apply the Claim above. Lemma 13 excludes the case $\delta_{n-2} = \delta_n = \delta_{n+2}$, so we have either $\delta_n = \delta_{n+4} \neq \delta_{n+2}$ or $\delta_n = \delta_{n-4} \neq \delta_{n-2}$. We treat the former case; the latter is similar (and may be reduced to the former by applying ρ to every string in question).

By Lemma 4, we have $\delta_{n+2} \ 2g \rightarrow \delta_n \ \delta_n$ but $\delta'_{n+2} \ 2g \rightarrow \delta'_n \ \delta'_{n+4}$, so if $\delta'_n \neq \delta_n$ we can't have $\delta'_{n+2} = \delta_{n+2}$.

But now the Claim above applies also to δ_{n+2} , and since $\delta_n, \delta_{n+4} \neq \delta_{n+2}$, we must have $\delta_{n-2} = \delta_{n+2} = \delta_{n+6}$.

Since $\delta_{n+2} \ 2g \rightarrow \delta_n \ \delta_n$ but $4g \neq 0$, by L5, $2\delta_{n+2} = 0$. Similarly, since $\delta_n \ 2g \rightarrow \delta_{n+2} \ \delta_{n+2}$, we have $2\delta_n = 0$.

But an application of A4 gives $2\delta_{n+1} \ 2g \rightarrow 2\delta_n \ 2\delta_{n+2} = 0 \ 0$, so $4\delta_{n+1} = 4g = 0$, a contradiction. This completes the proof that $+$ is associative. $\square$

7. 4-TORSION KUMMER STRUCTURES

In the previous section we have shown that every Kummer structure containing an element g with $4g \neq 0$ is the Kummer of a group. Here we turn to a different method to analyse the structure of a 4-torsion Kummer structure – i.e., one in which $4g = 0$ for all elements g . So throughout this section K shall be a 4-torsion Kummer structure.

To begin with, in section 3 we showed that $K[2]$, the 2-torsion of K , is a 2-torsion abelian group. For all $a \in K$, we have assumed $4a = 0$, or equivalently $2a \in K[2]$. So $2K$, the image of 2, is contained in $K[2]$, and it is easy to show (using L1 and A4) that $2K$ is a subgroup of $K[2]$.

If $a \in K$ and $x \in K[2]$, we can define $a + x$ by the rule $a x \rightarrow (a + x) (a + x)$.

Lemma 15. *This extends the group operation in $K[2]$ to an action of $K[2]$ on K , and if $a b \rightarrow c d$ then $a (b + x) \rightarrow (c + x) (d + x)$. For any $a \in K$, $x \in K[2]$, we have $2(a + x) = 2a$. If $a, b \in K$ with $2a = 2b$ then we can write $b = a + x$ for $x \in K[2]$, and x is unique modulo $2a$.*

Proof. That $+$ is a group action means that $a + 0 = a$ and $a + (x + y) = (a + x) + y$. The former follows from A1, while the latter and the statement $a b \rightarrow c d \Rightarrow a (b + x) \rightarrow (c + x) (d + x)$ follow from A3:

$$\begin{array}{ccc} \begin{array}{ccc} y & & y \\ a & x & \rightarrow (a+x) \end{array} & \begin{array}{ccc} y & & y \\ (a+x) & & (a+x) \end{array} & \begin{array}{ccc} x & & x \\ a & b & \rightarrow c \end{array} \\ \downarrow & \downarrow & \downarrow \\ \begin{array}{ccc} a & (x+y) & \rightarrow (a+x)+y \end{array} & \begin{array}{ccc} (a+x)+y & & (a+x)+y \end{array} & \begin{array}{ccc} a & (b+x) & \rightarrow (c+x) \end{array} \\ a & (x+y) & \rightarrow (a+x)+y \end{array} \quad \begin{array}{ccc} x & & x \\ a & (b+x) & \rightarrow (c+x) \end{array}$$

By A4 we have $2a 0 \rightarrow 2(a+x) 2(a+x)$, so $2(a+x) = 2a$. For a, b with $2a = 2b$, we know $2a 2b \rightarrow 0 0$, so if $a b \rightarrow x y$ then (by A4) $2x = 2y = 0$, and (by L2) $a x \rightarrow b \star$, so $b = a + x$. Also $b = a + y$, but (by L4) $x + y = 2a$, so $x \equiv y \pmod{2a}$. Any z with $b = a + z$ satisfies $a z \rightarrow b \star$, so $a b \rightarrow z \star$, so $z = x$ or y . $\square$

Thus if for each $x \in 2K$ we pick a representative $e_x \in K$ with $2e_x = x$, then every element of $a \in K$ can be written as $a = e_x + u$ with $x \in 2K$, $u \in K[2]$; given a , $x = 2a$ is unique and u unique modulo x .

Lemma 16. *Given a choice of $\{e_x\}$, there are elements $\varepsilon_{x,y} \in K[2]$ for each $x, y \in 2K$, such that the structure map of K is given by*

$$(e_x + u) (e_y + v) \rightarrow (e_{x+y} + u + v + \varepsilon_{x,y}) (e_{x+y} + u + v + \varepsilon_{x,y} + x).$$

$\varepsilon_{x,y}$ is well-defined modulo $\langle x, y \rangle$.

Proof. Let $e_x e_y \rightarrow a \star$. Then (by A4) $2e_x 2e_y \rightarrow 2a \star$. But $2e_x = x$ and $2e_y = y$, so $2a = x + y$. So $a = e_{x+y} + \varepsilon$ for some $\varepsilon \in K[2]$, defined (given the choice of a) modulo $\langle x + y \rangle$.

Now suppose $e_x e_y \rightarrow a b$. By L4, $a b \rightarrow x y$, so $a x \rightarrow b \star$, so $b = a + x = e_{x,y} + \varepsilon + x$. Thus choosing b instead of a changes ε by x , so overall the equation $e_x e_y \rightarrow (e_{x+y} + \varepsilon_{x,y}) \star$ defines $\varepsilon_{x,y}$ modulo $\langle x, y \rangle$.

Using this as the definition of $\varepsilon_{x,y}$, two applications of Lemma 15 give the full structure map of K as in the lemma. $\square$

A 4-torsion Kummer structure is thus determined up to isomorphism by the pair of groups $2K \subseteq K[2]$ and the pairing $(x, y) \in (2K)^2 \mapsto \varepsilon_{x,y} \in K[2]/\langle x, y \rangle$. But the pairing $\varepsilon_{\cdot, \cdot}$ is not canonical, since it depends on the choice of representatives $\{e_x\}$. We shall investigate its properties and try to find a standard form into which we can put $\varepsilon_{\cdot, \cdot}$. To begin with, we can take $e_0 = 0$, and from now on we shall do so. $\varepsilon_{x,y}$ is only defined modulo $\langle x, y \rangle$, so we take $\varepsilon_{x,y} = z$ to mean the same as $\varepsilon_{x,y} \equiv z \pmod{\langle x, y \rangle}$.

Lemma 17. *For any $x, y, z \in 2K$,*

$$\varepsilon_{x,y} + \varepsilon_{x+y,z} \equiv \varepsilon_{x,z} + \varepsilon_{x+z,y} \pmod{\langle x, y, z \rangle}.$$

Proof. Define $\delta \in K[2]$ by

$$\begin{array}{ccccc} & e_z & & e_z & e_z \\ e_y & e_x & \rightarrow & (e_{x+y} + \varepsilon_{x,y}) & \star \\ & \downarrow & & \downarrow & \downarrow \\ e_y & (e_{x+z} + \varepsilon_{x,z}) & \rightarrow & (e_{x+y+z} + \delta) & \star \\ e_y & \star & \rightarrow & \star & \star. \end{array}$$

The vertical arrow gives $\delta \equiv \varepsilon_{x,y} + \varepsilon_{x+y,z} \pmod{\langle x+y, z \rangle}$ and the horizontal arrow gives $\delta \equiv \varepsilon_{x,z} + \varepsilon_{x+z,y} \pmod{\langle x+z, y \rangle}$. Putting these together gives the lemma. $\square$

Lemma 18. $\varepsilon_{x,y} \in K[2]/\langle x, y \rangle$ depends only on the subgroup $\langle x, y \rangle \subseteq 2K$, not on the choice of generators x, y . If $\text{rank}\langle x, y \rangle < 2$ then $\varepsilon_{x,y} = 0$.

Proof. $A_2 \Rightarrow e_x e_x \rightarrow 0$. By definition, $e_x e_x \rightarrow (e_0 + \varepsilon_{x,x}) \star$, so $\varepsilon_{x,x} = e_0 = 0$.

Now let $z = x + y$: Lemma 17 becomes $\varepsilon_{x,y} + \varepsilon_{z,z} \equiv \varepsilon_{x,z} + \varepsilon_{y,y} \pmod{\langle x, y \rangle}$. Since $\varepsilon_{z,z} \equiv 0 \equiv \varepsilon_{y,y} \pmod{\langle x, y \rangle}$, we have $\varepsilon_{x,y} \equiv \varepsilon_{x,z} \pmod{\langle x, y \rangle}$; that is, $\varepsilon_{x,y} = \varepsilon_{x,z}$. This, together with the obvious symmetry $\varepsilon_{x,y} = \varepsilon_{y,x}$, shows that $\varepsilon_{x,y}$ depends only on $\langle x, y \rangle$ (since these two symmetries generate the group $GL_2(\mathbb{Z}/2)$).

If $\text{rank}\langle x, y \rangle < 2$ then $\langle x, y \rangle$ is generated by a single element z , and $\varepsilon_{x,y} = \varepsilon_{z,z} = 0$. $\square$

In the light of the last lemma, if $V = \langle x, y \rangle$ then we shall use the notation $\varepsilon_V = \varepsilon_{x,y}$.

Lemma 19. Let $\{x_1, \dots, x_n\}$ be a basis for a rank- n subgroup $X \subseteq 2K$. Call two elements of this subgroup disjoint if they are sums of disjoint subsets of the basis. Then we can choose $\{e_x\}$ such that when $a, b \in X \setminus 0$ are disjoint, $\varepsilon_{a,b} = 0$, and if $a, b, c \in X \setminus 0$ are disjoint, $\varepsilon_{a+b,a+c} \equiv 0 \pmod{\langle a, b, c \rangle}$.

Proof. We work by induction on n : for $n = 0$ there is nothing to prove, so let $X' = \langle x_1, \dots, x_{n-1} \rangle$ and suppose we have already chosen e_x for all $x \in X'$ satisfying the given conditions on ε_V for $V \subseteq X'$. Define $T = x_1 + \dots + x_n$ and $T' = T - x_n$.

Choose e_{x_n} arbitrarily (subject to $2e_{x_n} = x_n$). Then for each $a \in X' \setminus 0$, we can pick e_{a+x_n} such that $e_a e_{x_n} \rightarrow e_{a+x_n} \star$, so that $\varepsilon_{a,x_n} = 0$. There are two such choices, differing by x_n . If also $a \neq T'$, then by Lemma 17,

$$\varepsilon_{a,x_n} + \varepsilon_{a+x_n,T'-a} \equiv \varepsilon_{a,T'-a} + \varepsilon_{T',x_n} \pmod{\langle a, T' - a, x_n \rangle}.$$

By induction $\varepsilon_{a,T'-a} = 0$ and by the choices just made $\varepsilon_{a,x_n} = 0$ and $\varepsilon_{T',x_n} = 0$. So $\varepsilon_{a+x_n,T'-a} \equiv 0 \pmod{\langle a+x_n, T' - a, x_n \rangle}$. By adding x_n to e_{a+x_n} if necessary, we can make $\varepsilon_{a+x_n,T'-a} = 0$. So we have now $\varepsilon_{a,b} = 0$ whenever $a \in X'$ and $b = x_n$ or when $a + b = T$.

Now let $a, b \in X$ be disjoint. If $a, b \in X'$, we know $\varepsilon_{a,b} = 0$ by induction. So assume $a = a' + x_n$ with $a', b \in X'$. Lemma 17 says

$$\varepsilon_{a',x_n} + \varepsilon_{a,b} \equiv \varepsilon_{a',b} + \varepsilon_{a'+b,x_n} \pmod{\langle a, b, x_n \rangle}.$$

Three of these terms we know to be 0, leaving $\varepsilon_{a,b} \equiv 0 \pmod{\langle a, b, x_n \rangle}$. Also by Lemma 17,

$$\varepsilon_{b,a} + \varepsilon_{b+a,T-a-b} \equiv \varepsilon_{b,T-a-b} + \varepsilon_{T-a,a} \pmod{\langle a, b, T \rangle}.$$

Again three of these terms we know to be 0, leaving $\varepsilon_{b,a} = \varepsilon_{a,b} \equiv 0 \pmod{\langle a, b, T \rangle}$. If $T \notin \langle a, b, x_n \rangle$, we can put these together to get $\varepsilon_{a,b} = 0$. But if $T \in \langle a, b, x_n \rangle = \langle a', b, x_n \rangle$ then $T = a' + b + x_n = a + b$, since a', b , and x_n are disjoint, and in this case $\varepsilon_{a,b} = 0$ from the choices made in the previous paragraph.

Finally let $a, b, c \in X$ be disjoint and non-zero. Once more Lemma 17 says

$$\varepsilon_{a,b} + \varepsilon_{a+b,a+c} \equiv \varepsilon_{a,a+c} + \varepsilon_{c,b} \pmod{\langle a, b, c \rangle},$$

and $\varepsilon_{a,b} = 0$, $\varepsilon_{a,a+c} = \varepsilon_{a,c} = 0$, $\varepsilon_{c,b} = 0$, leaving $\varepsilon_{a+b,a+c} \equiv 0 \pmod{\langle a, b, c \rangle}$ as required. $\square$

Definition. Let $X = \langle x, y, z \rangle \subseteq 2K$ be a rank-3 subgroup. Define $\text{ind}_X = 0$ if there is a choice of $\{e_x\}$ such that $\varepsilon_V = 0$ for all rank-2 $V \subset X$; otherwise, define $\text{ind}_X = 1$.

Lemma 20. Let $X = \langle a, b, c \rangle$ be rank 3, and let $\{e_x\}$ be chosen as allowed by Lemma 19 so that $\varepsilon_V = 0$ for all $V \subset X$ except possibly for $V = U := \langle a + b, a + c \rangle$. Then $\varepsilon_U = 0 \Leftrightarrow \text{ind}_X = 0$.

Proof. We know $\varepsilon_U \equiv 0 \pmod{X}$. If $\varepsilon_U = 0$ then we have $\text{ind}_X = 0$ by definition. So all we need to prove is that if $\text{ind}_X = 0$ then $\varepsilon_U = 0$.

Suppose $\text{ind}_X = 0$. Then there is another choice of representatives $\{e'_x\}$ giving $\varepsilon'_V = 0$ for all $V \subset X$. By Lemma 15, each $e'_x = e_x + \delta_x$ for some $\delta_x \in K[2]$. So $e'_x e'_y \rightarrow (e'_{x+y} + \varepsilon_{x,y} + \delta_x + \delta_y + \delta_{x+y}) \star$; but for all $x, y \in X$, we know $e'_x e'_y \rightarrow e'_{x+y} \star$, so

$$\varepsilon_{x,y} \equiv \delta_x + \delta_y + \delta_{x+y} \pmod{\langle x, y \rangle}.$$

Set $\Delta := \delta_a + \delta_b + \delta_c + \delta_{a+b+c}$. Since $\varepsilon_V = 0$ for all $U \neq V \subset X$, we have

$$\delta_{a+b} + \delta_a + \delta_b \equiv 0 \pmod{\langle a, b \rangle}$$

$$\delta_{a+b} + \delta_c + \delta_{a+b+c} \equiv 0 \pmod{\langle a + b, c \rangle},$$

and, adding these together, we can see that $\Delta \in X$ and

$$\begin{aligned} \Delta \in \langle a + b, c \rangle &\Leftrightarrow \delta_{a+b} + \delta_a + \delta_b \equiv 0 \pmod{\langle a + b \rangle} \\ &\Leftrightarrow \delta_{a+b} + \delta_a + \delta_b \in U. \end{aligned}$$

Analogous statements hold for other permutations of $\{a, b, c\}$.

Now it is easy to check that every element of X is in an even number of the subgroups $\langle a + b, c \rangle$, $\langle a + c, b \rangle$, and $\langle b + c, a \rangle$. So an even number out of $(\delta_{a+b} + \delta_a + \delta_b)$, $(\delta_{a+c} + \delta_a + \delta_c)$, and $(\delta_{b+c} + \delta_b + \delta_c)$ are in U , so their sum is in U . But their sum is

$$\delta_{a+b} + \delta_{a+c} + \delta_{b+c} \equiv \varepsilon_U \pmod{U},$$

so $\varepsilon_U = 0$, as required. $\square$

Lemma 21. If X and Y are two rank-3 subgroups of $2K$ then $\text{ind}_X = \text{ind}_Y$.

Proof. It is enough to show this when $\text{rank}(X \cap Y) = 2$, since any two rank-3 subgroups can be connected by a chain of subgroups with each 2 consecutive members of the chain meeting having rank-2 intersection.

So we may assume $X = \langle a, b, c \rangle$ and $Y = \langle b, c, d \rangle$, and choose $\{e_x\}$ by Lemma 19. Let $U := \langle a + b, a + c, a + d \rangle$, the group of even sums of $\{a, b, c, d\}$. By Lemma 17,

$$\varepsilon_{a+b,a+c} + \varepsilon_{b+c,c+d} \equiv \varepsilon_{a+b,c+d} + \varepsilon_{a+b+c+d,b+c} \pmod{U}.$$

But according to Lemma 19, the two terms on the right-hand side of this are 0, while by Lemma 20, $\varepsilon_{a+b,a+c} \in U \Leftrightarrow \text{ind}_X = 0$ and $\varepsilon_{b+c,c+d} \in U \Leftrightarrow \text{ind}_Y = 0$. $\square$

So if $\text{rank } 2K \geq 3$ we can define $\text{ind}_K := \text{ind}_X$ for any rank-3 $X \subseteq K$. Finally we can classify 4-torsion Kummer structures:

Theorem 22. Let K be a 4-torsion, but not 2-torsion, Kummer structure.

$K \cong G/\pm 1$ for an abelian group G if and only if $\text{rank } 2K \leq 2$ or $\text{ind}_K = 1$.

$K \cong G/\iota$ for a twisted group (G, ι) if and only if $\text{rank } 2K \leq 2$ or $\text{ind}_K = 0$.

Proof. For the only-if parts, note that for $K \cong G/\pm 1$, a rank-3 $X \subseteq 2K$ is the image of a rank-3 subgroup of $2G$, which is $2\times$ a subgroup of G isomorphic to $(\mathbb{Z}/4)^3$. It is easy to check then that $\text{ind}_X = 1$. Similarly, if $K \cong G/\iota$, a rank-3 $X \subseteq 2K$ is the image of a rank-3 subgroup of $(1 + \iota)G$, which is $(1 + \iota)$ times a subgroup isomorphic to $((\mathbb{Z}/2)^2, \iota_2)^3$, where ι_2 exchanges the two factors of $(\mathbb{Z}/2)^2$. It is again easy to check that $\text{ind}_X = 0$.

For the if parts, we have shown that K is determined up to isomorphism by the two groups $2K \subseteq K[2]$ and a pairing $(x, y) \in (2K)^2 \mapsto \varepsilon_{x,y} \in K[2]/\langle x, y \rangle$. If K is finite, for any basis of $2K$ we can put $\varepsilon_{\cdot,\cdot}$ in standard form as in Lemma 19; then by Lemmas 20 and 21, the pairing $\varepsilon_{\cdot,\cdot}$ is entirely determined by ind_K . So a finite 4-torsion Kummer structure K is determined up to isomorphism by $\text{rank } 2K$, $\text{rank } K[2]$, and, if $\text{rank } 2K \geq 3$, ind_K .

But if $G = (\mathbb{Z}/4)^a \oplus (\mathbb{Z}/2)^b$ and $K = G/\pm 1$ then $\text{rank } 2K = a$, $\text{rank } K[2] = a + b$, and, if $a \geq 3$, $\text{ind}_K = 1$. And if $(G, \iota) = ((\mathbb{Z}/2)^2, \iota_2)^a \oplus (\mathbb{Z}/2, 1)^b$ and $K = G/\iota$ then $\text{rank } 2K = a$, $\text{rank } K[2] = a + b$, and, if $a \geq 3$, $\text{ind}_K = 0$.

This proves the theorem for finite K . For infinite K , we can see from Theorem 5 that the property of being isomorphic to the Kummer of a group, and the property of being isomorphic to a twisted Kummer, are each equivalent to a collection of statements that depend only on the structure of finitely generated substructures (for example, the statement that $\alpha + (\beta + \gamma) = (\alpha + \beta) + \gamma$ in K_g depends only on the substructure generated by $\{\alpha_0, \beta_0, \gamma_0, g\}$). A finitely generated substructure L of a 4-torsion Kummer structure K is finite, and if one of the properties $\text{rank } 2K \leq 2$ or $\text{ind}_K = i$ holds, the same property holds for every sub-Kummer-structure of K , so the theorem can be extended from finite K to arbitrary K . $\square$

8. SUMMARY

Collecting together Theorems 5, 14, and 22, and the Remark after Lemma 2, we have the following classification of Kummer structures:

Theorem 23. *Every Kummer structure is either the Kummer of an abelian group or a twisted Kummer. If $K \cong G/\pm 1$ then G is determined up to isomorphism by K . If $K \cong G/\iota$ then (G, ι) is determined up to isomorphism by K . The only Kummers that are also twisted Kummers are $G/\pm 1$ where either G is 2-torsion or G is 4-torsion and $\text{rank } 2G \leq 2$, which are isomorphic to G/ι where $\text{rank}(1 + \iota)G \leq 2$. $\square$*

We can therefore characterise Kummers of groups by adding to A1–A4 the statement that K has no substructure isomorphic to the unique twisted Kummer L with $\text{rank } 2L = \text{rank } L[2] = 3$, which is $(\mathbb{Z}/2)^6/\iota$, where the involution ι acts on a basis $\{x_0, \dots, x_5\}$ by $x_i \mapsto x_{(i+3) \bmod 6}$.

Remark. In fact, G or (G, ι) is determined by K in a constructive and computable way. For 2-torsion K this is trivial, while for non-2-torsion K Theorem 5 shows that the elements of G can be represented as strings, and Lemma 3 shows that a string can be represented by a pair of elements of K . The string $\gamma = \alpha + \beta$ or $\gamma = \alpha \oplus \beta$ of Section 4 is determined up to finite ambiguity from $\alpha_0 \beta_0 \rightarrow \gamma_0 \star$, and the proof of Lemma 12 shows that this ambiguity can be removed by extending the strings to a sufficient (finite) length and considering all the conditions on γ_0 and γ_1 that then arise.